Junior Quiz

| 1 | How long is the average gestation length for beef cattle? | | | | |
|--|---|---|-----------------------|-------------------------------|--|
| 1. | - | | c | 292 dave | |
| | | days | <mark>ر.</mark> ما | 283 days | |
| h | b. 145 days d. 320 days | | | | |
| A cow's nutritional requirements would be highest durin Maintenance | | | | | |
| | | | C. | 107 | |
| 2 | | k lactation | a. | Breeding season | |
| 3. | | s classified as: | _ | Duchistic | |
| | | tein supplement | - | Probiotic | |
| | | ophore | d. | Antibiotic | |
| 4. | | acceptable amount of back fat for a market la | | 0.45 | |
| | | 5 inches | с. | 0.45 inches | |
| _ | | 5 inches | d. | A and B | |
| 5. | 5. Which of the following species is a ruminant? | | | | |
| | a. Swi | | <mark>C.</mark> | Cattle | |
| _ | b. Hor | | d. | | |
| 6. | 6. Intramuscular fat in a cut of meat is known as which of the following? | | | | |
| | | l fat | с. | Back fat | |
| | | rbling | d. | | |
| 7. | | | | | |
| | - | ly gain (ADG) of the steer? | | | |
| | | pounds | с. | 2.0 pounds | |
| | b. 1.5 | | d. | 3.5 pounds | |
| 8. | 8. Which of the following are USDA cattle quality grades? | | | | |
| | | ne, Choice, Select | с. | , , , , , | |
| | - | s, C, D, E | | Prime, Choice, Good | |
| 9. | . Hardware disease is typically associated with which compartment of the stomach? | | | | |
| | a. Rur | nen | с. | Omasum | |
| | | iculum | d. | Abomasum | |
| 10. Which animal is <i>most</i> likely to experience ketosis? | | | | | |
| | a. Ab | eef cow pregnant with a | <mark>C.</mark> | A ewe carrying multiple lambs | |
| | sing | le fetus | d. | All of the above | |
| | b. Ad | airy cow in positive energy | | | |
| | bala | ance | | | |
| 11. A calf deficient in selenium would most likely experience which of the following? | | | | | |
| | a. Ent | erotoxemia | с. | Polio | |
| | <mark>b. Wh</mark> | ite muscle disease | d. | Mastitis | |
| 12. | 12. Bovine spongiform encephalopathy, or BSE, is a condition caused by: | | | | |
| | a. Pro | tein deficiency | c. | Prions | |
| | b. Rin | gworm | d. | Barber Poll worm | |
| | | | _ | | |

13. A USDA yield grade of ______ would indicate an extremely fat, lightly muscled carcass.

| a. 1 | c. 4 | | | | |
|---|---|--|--|--|--|
| b. 2 | d. 5 | | | | |
| 14. This breed of sheep is noted for its parasite resistance | | | | | |
| a. Dorset | c. Suffolk | | | | |
| b. Merino | <mark>d. Katahdin</mark> | | | | |
| 15. If a cow has a body condition score of 2, she is best described as: | | | | | |
| a. Ideal | c. Fat | | | | |
| <mark>b. Thin</mark> | d. Obese | | | | |
| 16. A cow suffering from grass tetany is deficient in which mineral? | | | | | |
| a. Calcium | c. Magnesium | | | | |
| b. Manganese | d. Copper | | | | |
| 17. A sheep with which of the following genotypes at codon 171 would be susceptible to Scapie? | | | | | |
| a. QR | c. QQ | | | | |
| b. RR | d. RR | | | | |
| 18. Which of these primal cuts represent the lowest percent | age of carcass weight? | | | | |
| a. Shoulder (Butt) | c. Ham | | | | |
| b. Loin | d. Side | | | | |
| 19. Your current sow population consists primarily of Yorkshire and Landrace genetics. Which of the | | | | | |
| following breeds could be utilized to increase lean growth and improve feed efficiency of the | | | | | |
| market hogs produced? | | | | | |
| a. Berkshire | c. Chester White | | | | |
| <mark>b. Hampshire</mark> | d. Hereford | | | | |
| 20. Your ewe was marked on October 1. If she did not become | ne pregnant, what day would you | | | | |
| expect her to be in heat again? | | | | | |
| a. October 7 | c. October 18 | | | | |
| b. October 28 | d. November 1 | | | | |
| 21. What hormone is released by a CIDR? | | | | | |
| a. Progesterone | c. Estradiol | | | | |
| b. Prostaglandin | d. Oxytocin | | | | |
| 22. What mineral is toxic to sheep but can be fed to other liv | | | | | |
| a. Zinc b. Iron | <mark>c. Copper</mark> d. Selenium | | | | |
| | | | | | |
| 23. What is a defect called when a sheep's bottom jaw extera. Parrot mouth | | | | | |
| a. Parrot mouth b. Broken mouth | c. Spreader <mark>d. Monkey mouth</mark> | | | | |
| 24. Which parasite is prevalent in the Southeastern US and c | | | | | |
| a. Coccidia | c. Ticks | | | | |
| | d. Brown Stomach Worm | | | | |
| b. Barber Pole Worm 25. What is the average dressing percentage of market hogs | | | | | |
| | с. 52% | | | | |
| | d. 90% | | | | |
| b. 62.5% | u. 30/0 | | | | |